COURSE GUIDE

CHM 318 NATURAL PRODUCT CHEMISTRY I

Course Team Dr. I. A. Ogunwande (Course Writer) -

Lagos, Nigeria

Prof. Kehinde Obidairo (Course Editor)- NOUN Prof. Dimas. Kubmarawa (Course Reviewer) -

Modibbo Adama University of Tech. Yola



NATIONAL OPEN UNIVERSITY OF NIGERIA

© 2021 by NOUN Press
National Open University of Nigeria
Headquarters
University Village
Plot 91, Cadastral Zone
Nnamdi Azikiwe Expressway
Jabi, Abuja

Lagos Office 14/16 Ahmadu Bello Way Victoria Island, Lagos

e-mail: <u>centralinfo@nou.edu.ng</u>

URL: www.nou.edu.ng

All rights reserved. No part of this book may be reproduced, in any form or by any means, without permission in writing from the publisher.

Printed: 2021

ISBN: 978-978-058-079-7

Revised in 2021

CONTENTS	PAGE
Introduction	iv
What you will Learn in this Course	iv
Course Aim	iv
Course Objectives	iv
Working through this Course	V
Study Units	V
Presentation Schedule	vi
Assessment	vi
Tutor-Marked Assignment (TMA)	vi
Final Examination and Grading	vii
Course Marking Scheme	vii
Facilitators/Tutors and Tutorials	vii
Summary	viii

INTRODUCTION

Natural Product Chemistry 1 is a semester course which is part of the courses designated as part of requirements for the award of B.Sc. Chemistry

WHAT YOU WILL LEARN IN THIS COURSE

The course consists of units and a course guide. The course guide is primarily a simple description of what the course is made up of, the materials to be use and how you can work with these materials. In addition, it advocates some general guidelines for the time you are likely to spend on each unit of the course in order to complete it successfully.

Also, you find in it your Tutor Marked Assignment which will be made available in the assignment file. Moreover, there are regular tutorial questions and classes that will assist in the better understanding of the course. You are advised to make yourselves available during the tutorial classes.

COURSE AIM

The aim of the course is to provide a basic understanding of the basic nature of chemical materials and the emerging trend. In addition, it seeks to address the massive drive to understand these materials and improve their properties in order to meet material requirements.

COURSE OBJECTIVES

To achieve the set aim, the course has a set of objectives. Each of the unit contained in a specific module has stated objectives which are included at the beginning of the particular unit. A clearer understanding of each of the objectives is a prerequisite for the better comprehension of the contents of the unit. It is highly essential to reflect, as you work through each unit, on the objectives.

The main objectives of the course are listed below. By meeting these objectives, you should have achieved the aims for which the course has been studied. At the end of this course, you should be able to:

- Know what natural products are
- Understand the different types of natural products
- Give various examples of natural products
- Describe chemical properties of natural products
- State the applications of these natural products

WORKING THROUGH THIS COURSE

In order to be able to successfully complete this course, you are required to carefully study each unit along with recommended textbooks and other materials that may be provided by the National Open University. You may also need to exploit other e-reading such internet for further useful information on the course.

Each unit contains self-assessment exercise and at certain points in the course you would be required to submit assignment for grading and recording purposes. You are also to participate in the final examination at the end of the course. It is recommended that you devote an abundant time for reading and comprehension. It is highly necessary that you avail yourselves the opportunity of attending the tutorial sessions where you will be able to compare your understanding of the course contents with your colleagues.

The Course Materials

The main components of this course are:

- 1. The Course Guide
- 2. Study Units
- 3. Self Assessment Exercise
- 4. Self-Assessment Questions
- 5. Further Readings

STUDY UNITS

The study units in this course are as follows:

Module 1 Terpenes

Unit 1 Biosynthesis of Terpenes Unit 2 Classification of Terpenes

Module 2 Steroids

Unit 1 Nature of Steroids

Module 3 Alkaloids

Unit 1 Nature of Alkaloids

Unit 2

Classification of AlkaloidsThe first unit addresses the nature of terpenes. It focuses on history, biosynthesis and importance of terpenes. The second unit described in details the different of terpenes and their examples. The third unit is primarily concerned with the synthesis and structures of steroids. The fourth unit addresses the properties, distribution, extraction and application of alkaloids. The fifth unit deals with the diversity of alkaloids and their specific importance.

Each of the unit is made up of one or two weeks' work consisting of introduction, objectives, reading materials, self assessment exercise, conclusion, summary and Tutor marked Assignment (TMA), suggestion for further reading and source materials. The unit directs you to work on exercises related to the required reading. Together with the TMAs, they are meant to test your basic understanding and comprehension of the course materials, which is a prerequisite for the achieving the stated aims and objectives of the course.

PRESENTATION SCHEDULE

The course materials have important dates for the timely completion and submission of your TMAs and tutorial lessons. You are vividly reminded of the need to promptly submit answers to tutorials and assignments as at when due.

ASSESSMENT

The course assessment consists of three aspects namely the self assessment exercise, the tutor marked assignment and the written examination/end of course examination.

It is essential that you attempt all exercises and assignments and submit appropriately to the course facilitator for grading. Let your answers be concise and as accurate as possible. You are expected to consult other material course in addition to your course materials in order to be able to present accurate answers to the questions. Kindly note that the tutor marked assignment covers only 30% of the total marked for the course.

TUTOR MARKED ASSIGNMENT (TMA)

The TMA is a continuous assessment component of your course. It accounts for 30% of the total score. You will be given five (5) TMAs to

answer. Three of these must be answered before you are allowed to sit for the end of the course examination. The TMAs will be given to you by your facilitator and returned after you have done the assignment. Note that these assignments are already contained in the assignment file to be given to you. You may do yourself good by reading and researching well before you attempt to answer the questions.

You are warned to submit these assignments to the facilitator at the stipulated time as could be seen in the assignment file. However, if for any reason you are unable to meet the deadline, you are highly required to intimate the facilitator of your problem before the due date and seek for an extension which may be granted or not.

FINAL EXAMINATION AND GRADING

The end of the course examination for Natural Product Chemistry 1 will be for about 3 hours with maximum score value of 70% of the total course work. The examination will be made up of questions which normally reflect on what you have learnt in the course materials/further reading. In addition, they may be prototype of the self assessment exercises and the TMAs. The end of the course examination is intended to cover information from all parts of the course.

Avail yourself the opportunity of the time-lag between the completion of the course content and the beginning of the examination to revise as much as possible the whole course materials, the exercise and assignments.

COURSE MARKING SCHEME

Assignment	Marks
	Five assignments, best three marks of
Assignments 1-5	the
	five count at 10% each i.e. 30% of the
	course marks
End of course Examination	70% of overall course marks
Total	100% of the course materials

FACILITATORS/TUTORS AND TUTORIALS

There are 17 hours of tutorials provided in support of this course. You will be informed appropriately of the name, telephone number and email address of your facilitator. In addition, the time, dates and location of the tutorial lessons will be communicated beforehand. You are required to mail or submit your Tutor Marked Assignment to your facilitator, at least two working days, before the schedule date. Note that

all the submitted assignments will be duly marked by the facilitator with further comments that can improve on your performances. The facilitator will from time to time takes track record of your comprehension, progress and difficulty in the course.

Be kind enough to attend tutorial lessons at the fixed appointment. It is probably the only avenue to meet face to face and discuss with you facilitator. There, you will be able to ask question or seek clarification on seemingly grey area in the course material. You may as well have prepared questions and comments for your facilitator before the due date. An active participation during the tutorial lessons will be an added advantage to boost confidence level.

In case any of the situations listed below arises, do not hesitate to intimate your facilitator using his or her telephone number or via e-mail address;

- You do not understand any part of the study or the assigned readings
- You are not skill enough to attempt the self-assessment exercise the questions in the TMAs are not clearly understood

SUMMARY

Natural Product chemistry is a course which is intended to provide students with the nature and classes of natural products. Upon completion of this course, you will be highly equipped to answer questions below and related ones:

- . What do natural products represent?
- What are the different type, chemical nature and features of natural products?
- Know different classification of natural products
- . Identify the examples of terpenes, steroids and alkaloids.
- Discuss the application of terpenes, steroids and alkaloids.

Accept my best wishes in the course and I do hope that you benefit considerably from its application.

MAIN COURSE

CONTENT	TS .	PAGE
Module 1		1
Unit 1 Unit 2	Biosynthesis of Terpenes	1 13
Module 2		25
Unit 1	Nature of Steroids	25
Module 3		36
Unit 1	Nature of Alkaloids	36

MODULE 1 TERPENES

UNIT 1 BIOSYNTHESIS OF TERPENES

CONTENTS

- 1.0 Introduction
- 2.0 Objectives
- 3.0 Definition of Terpenes
 - 3.1 History of Terpenes
 - 3.2 Biosynthesis of Terpenes
 - 3.2.1 Mevalonic acid pathway
 - 3.2.2 MEP/DOXP pathway
 - 3.3 Summarized classification of Terpenes
 - 3.4 Importance of Terpenes
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 References/Further Reading

1.0 INTRODUCTION

During the 19th century, chemical works on turpentine led to name "terpene" the hydrocarbons with the general formula $C_{10}H_{16}$ found in complex plant product. These terpenes are frequently found in plant essential oils which contain the "*Quinta essentia*", the plant fragrance. They are universally present in small amounts in living organisms, where they play numerous vital roles in plant physiology as well as important functions in all cellular membranes. On the other hand, they are also accumulated in many cases, and it is shown that the extraordinary variety they then display can be due to ecological factors playing an evolutionary role.

2.0 OBJECTIVES

By the end of this unit, students should be able to:

- define terpenes and terpenoids
- understand the history behind terpenes
- describe the biosynthesis of terpenes
- state the form in terpenes occur naturally

3.0 Definition of Terpenes and Terpenoids

The name "terpene" is derived from the word "turpentine". Terpenes may be defined as a group of molecules whose structure is based on a various but definite number of isoprene units (methylbuta-1,3-diene, named hemiterpene, with 5 carbon atoms). When terpenes are modified chemically, such as by oxidation or rearrangement of the carbon skeleton, the resulting compounds are generally referred to as *terpenoids*.

Terpenoids (or isoprenoids), a subclass of the prenyllipids (terpenes, prenylquinones, and sterols), represent the oldest group of small molecular products synthesized by plants and are probably the most widespread group of natural products. Terpenoids can be described as modified terpenes, where methyl groups are moved or removed, or oxygen atoms added. Inversely, some authors use the term "terpenes" more broadly, to include the terpenoids.

3.1 History of Terpenes

Terpenes history spans various civilizations. As they are largely found in essential oils, they were used in the Ancient Egypt for various religious aims. Camphor was introduced in Europe from the East by the Arabs around the 11th century. The process of obtaining plant essential oils by fatty extraction was known by the early Middle Ages. In the 12th century, Arnaud de Villanos described distillation of oils from rosemary and sage. He made an "oleum mirabile" from oils of turpentine and rosemary.

It is noticeable that some 60 oils were described in the Nuremberg edition of "*Dispensatorium valerii cordi*" written in 1592. Analyses of oils of turpentine were made in 1818 by JJ Houston de la Billardière. Dumas proposed in 1866 the name "terpene", derived from turpentine, instead of camphor for crystalline oxygenated substances extracted from

essential oils. In 1887, Wallach O, proposed that one isoprenic unit of 5 carbon atoms (C₅H₈) is always present in the molecule of terpenes.

The structure of camphor was established by Bredt in 1893, that of pinene by Wagner in 1894 and that of citral by Tiemann in 1895. - Carotene was isolated in 1837 by Wackenrodder from carrots, and its correct molecular form was determined in 1907 by Willstätter. The period since 1945 has seen an extensive explosion in natural product chemistry due to the advent of chromatographic and spectroscopic techniques.

-Carotene

The discovery of the isoprene unit is the basis of the concept of the "isoprenic rule" edicted in 1953 by Ruzicka and completed by Lynen. Mevalonic acid was shown in 1956 to be a biosynthetic precursor of cholesterol and later, its incorporation into a number of terpenoids has been demonstrated. Actually, an increasing number of terpenoids are described in the plant kingdom and many of them were shown to have important biological activities. Thus, several sesquiterpenes and diterpenes have antibiotic properties, some sesquiterpenes and diterpenes are insect and plant hormones, respectively.



Mevalonic acid

3.2 Biosynthesis of Terpenes

Terpenoids are extraordinarily diverse but they all originate through the condensation of the universal phosphorylated derivative of hemiterpene, isopentenyl diphosphate (IPP) and dimethylallyl diphosphate (DMAPP) giving geranyl pyrophosphate (GPP).

In higher plants, IPP is derived from the classic mevalonic acid pathway in the cytosol but from the methylerythritol phosphate pathway in plastids. It is generally accepted that the cytosolic pool of IPP serves as a precursor of sesquiterpenes, triterpenes, sterols and polyterpenes whereas the plastid pool of IPP provides the precursors of mono-, diand tetraterpenes.

There are two metabolic pathways of creating terpenoids: Mevalonic acid pathway and MEP/DOXP pathway.

3.2.1 Mevalonic acid pathway

Many organisms manufacture terpenoids through the HMG-CoA reductase pathway, the pathway that also produces cholesterol. The reactions take place in the cytosol. The pathway was discovered in the 1950s.

Formation of Mevalonic Acid from Acetyl Units

3.2.2 MEP/DOXP pathway

The 2-*C*-methyl-D-erythritol 4-phosphate/1-deoxy-D-xylulose 5-phosphate pathway (MEP/DOXP pathway), also known as non-mevalonate pathway or mevalonic acid-independent pathway, takes place in the plastids of plants and apicomplexan protozoa, as well as in many bacteria. It was discovered in the late 1980s.

Pyruvate and glyceraldehyde 3-phosphate are converted by DOXP synthase (Dxs) to 1-deoxy-D-xylulose 5-phosphate, and by DOXP reductase (Dxr, IspC) to 2-C-methyl-D-erythritol 4-phosphate (MEP). The subsequent three reaction steps catalyzed by 4-diphosphocytidyl-2-C-methyl-D-erythritol synthase (YgbP, IspD), 4-diphosphocytidyl-2-Cmethyl-D-erythritol kinase (YchB, IspE), and 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase (YgbB, IspF) mediate the formation of 2-*C*-methyl-D-erythritol 2,4-cyclopyrophosphate (MEcPP). **MEcPP** converted (*E*)-4-hydroxy-3-methyl-but-2-enyl to pyrophosphate (HMB-PP) by HMB-PP synthase (GcpE, IspG), and HMB-PP is converted to isopentenyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP) by HMB-PP reductase (LytB, IspH).

IPP and DMAPP are the end-products in either pathway, and are the precursors of isoprene, monoterpenoids (10-carbon), diterpenoids (20-carbon), carotenoids (40-carbon), chlorophylls, and plastoquinone-9 (45-carbon). Synthesis of all higher terpenoids proceeds via formation of geranyl pyrophosphate (GPP), farnesyl pyrophosphate (FPP), and geranylgeranyl pyrophosphate (GGPP).

Although both pathways, MVA and MEP, are mutually exclusive in most organisms, interactions between them have been reported in plants and few bacteria species.

Organism Pathways

Bacteria	MVA or MEP			
Archaea			MVA	
Green Algae	MEP			
		Plants	MVA MEP	and
Animals			MVA	
Fungi			MVA	

3.3 Summarized Classification of Terpenes

A rational classification of the terpenes has been established based upon the number of isoprene (or isopentane) units incorporated in the basic molecular skeleton:

	Terpenes	Isoprene units	Carbon atoms
1	Monoterpenes	2	10
2	Sesquiterpenes	3	15
3	Diterpenes	4	20
4	Sesterpenes	5	25
5	Triterpenes	6	30
6	Carotenoids	8	40
7	Rubber	> 100	> 500

Mono-, sesqui-, di-, and sesterpenes contain the isoprene units linked in a head to tail fashion. The triterpenes and carotenoids (tetraterpenes) contain two C15 and C20 units respectively linked head to head. Many terpenes are hydrocarbons, but oxygen-containing compounds such as alcohols, aldehydes or ketones are also found. These derivatives are frequently named terpenoids. Mono- and sesquiterpenes are the chief constituents of the essential oils while the other terpenes are constituents of balsams, resins, waxes, and rubber.

3.4 Importance of Terpenes

The terpenoids, sometimes called *isoprenoids*, are a large and diverse class of naturally-occurring organic chemicals similar to terpenes, derived from five-carbon isoprene units assembled and modified in thousands of ways. Most are multicyclic structures that differ from one another not only in functional groups but also in their basic carbon skeletons. These lipids can be found in all classes of living things, and are the largest group of natural products.

Oleoresin is a roughly equal mixture of turpentine (85% C10-monoterpenes and 15% C15-sesquiterpenes) and resin (C20-diterpene) that acts in many conifer species to seal wounds and is toxic to both invading insects and their pathogenic. A number of inducible terpenoid defensive compounds (*phytoalexins*) from angiosperm species are well. These include both sesquiterpenoid and diterpenoid types.

Isoprenoid units are also found within the framework of other natural molecules. Thus, indole alkaloids, several quinones (vitamin K), alcohols (vitamin E, vitamin A formed from -carotene), phenols, isoprenoid alcohols (also known as terpenols or polyprenols) also contain terpenoid fragments. The origin of the ubiquitous isoprene unit and its conversion into various compounds has been extensively studied.

VITAMINS K

VITAMIN E

Tocopherol


VITAMIN A

Retinol

Terpenes are a large and varied class of organic compounds, produced primarily by a wide variety of plants, particularly conifers, though also by some insects such as termites or swallowtail butterflies, which emit terpene from their osmeterium. They are the major components of resin, and of turpentine produced from resin. In addition to their roles as end-products in many organisms, terpenes are major biosynthetic building blocks within nearly every living creature.

Terpenes and terpenoids are the primary constituents of the essential oils of many types of plants and flowers. Essential oils are used widely as natural flavor additives for food, as fragrances in perfumery, and in traditional and alternative medicines such as aromatherapy. Synthetic variations and derivatives of natural terpenes and terpenoids also greatly expand the variety of aromas used in perfumery and flavors used in food additives. Vitamin A is an example of a terpene.

The aroma and flavor of hops, highly desirable in some beers, comes from terpenes. The following terpenes in hops namely: myrcene, b-pinene, b-caryophyllene, and a-humulene are found in large quantities.




- Pinene


humulene

Plant terpenoids are used extensively for their aromatic qualities. They play a role in traditional herbal remedies and are under investigation for antibacterial, antineoplastic, and other pharmaceutical functions. Terpenoids contribute to the scent of eucalyptus, the flavors of cinnamon, cloves, and ginger, and the color of yellow flowers. Well-known terpenoids include citral, menthol, camphor, Salvinorin A in the plant *Salvia divinorum*, and the cannabinoids found in *Cannabis*.


Cetral B (Neral)



Menthol



R- and S-Camphor



Salvinorin A

The steroids and sterols in animals are biologically produced from terpenoid precursors. Sometimes terpenoids are added to proteins, e.g., to enhance their attachment to the cell membrane; this is known as isoprenylation.

SELF-ASSESSMENT EXERCISE

- i. What are terpenes and terpenoids?
- ii. Using appropriate scheme, describe the biosynthesis of terpenes.
- iii. Highlight the importance of terpenes and terpenoids

4.0 CONCLUSION

The terpenoids sometimes called *isoprenoids*, are a large and diverse class of naturally-occurring organic chemicals similar to terpenes, derived from five-carbon isoprene units assembled and modified in thousands of ways. Most are multicyclic structures that differ from one another not only in functional groups but also in their basic carbon skeletons. These lipids can be found in all classes of living things, and are the largest group of natural products.

5.0 SUMMARY

In summary, we have learnt that

- Terpenoids may be defined as a group of molecules whose structure is based on a various but definite number of isoprene units.
- The scientist that proposed the name 'terpene' was Dumas in 1866 by deriving it from turpentine
- Terpenes can be synthesized from two pathway; mevalonic acid pathway and the non-mevalonate pathway.
- Terpenes are a large and varied class of organic compounds, produced primarily by a wide variety of plants and some animals like insects

6.0 TUTOR-MARKED ASSIGNMENT

- 1. Distinguish between terpenes and terpenoids
- 2. Enumerate the importance of terpenes
- 3. State five examples of terpenes
- 4. Give example of monoterpenes formed by 'head to head and head to tail' orientation

7.0 REFERENCES/FURTHER READING

- Tadhg P. Begley (2009). *Encyclopedia of Chemical Biology*. Wiley. ISBN 978-0-471-75477-0.
- Paul M Dewick (2002). Medicinal Natural Products. A Biosynthetic Approach. Second Edition.

Wiley. ISBN 0471496405.

Patrick GL (2013). "Medical Folklore". An introduction to medicinal chemistry (Fifth ed.). Oxford: Oxford University Press. ISBN 978-0-19-969739-7.

UNIT 2 CLASSIFICATION OF TERPENES

CONTENTS

- 1.0 Introduction
- 2.0 Objectives
- 3.0 Monoterpenes
 - 3.0.1 Acyclic monoterpenes
 - 3.0.2 Monocyclic monoterpenes
 - 3.0.3 Bicyclic monoterpenes
 - 3.1 Sesquiterpenes
 - 3.1.1 Acyclic sesquiterpenes
 - 3.1.2 Cyclic sesquiterpenes
 - 3.2 Diterpenes
 - 3.3 Sesterpenes
 - 3.4 Triterpenes
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 References/Further Reading

1.0 INTRODUCTION

A rational classification of the terpenes has been established based upon the number of isoprene (or isopentane) units incorporated in the basic molecular skeleton:

	Terpenes	Isoprene	Carbon
		Units	atoms
1	Monoterpenes	2	10
2	Sesquiterpenes	3	15
3	Diterpenes	4	20
4	Sesterpenes	5	25
5	Triterpenes	6	30
6	Carotenoids	8	40
7	Rubber	> 100	> 500

Mono-, sesqui-, di-, and sesterpenes contain the isoprene units linked in a head to tail fashion. The triterpenes and carotenoids (tetraterpenes) contain two C15 and C20 units respectively linked head to head.

2.0 OBJECTIVES

By the end of this unit, students should be able to:


- understand the various classes of terpenes
- know the distinguishing features of all classes of terpenes
- give examples of each classes of terpenes

3.0 MONOTERPENES

They are the terpenes that have been known for several centuries as components of the fragrant oils obtained from leaves, flowers and fruits. Monoterpenes, with sesquiterpenes, are the main constituents of essential oils. While a few, such as camphor, occur in a near pure form, most occur as complex mixtures, often of isomers difficult to separate. These essential oils have numerous actions, such as allelochemical functions between plants and between plants and predators. A role in wound healing was also observed.

3.0.1 Acyclic Monoterpenes


Among natural molecules, the followings are well known and have several structural isomers.



Defensive role of simple terpenes have been demonstrated as for more complex compounds. Ocimene and linalool (with farnesene) were shown to be produced by *de novo* biosynthesis in plants damaged by insect herbivores. These compounds likely mediate the interaction between herbivores and their natural enemies, attracted By terpenes.


3.0.2 Monocyclic Monoterpenes

They are derived from cyclohexane with an isopropyl substituent. The most typical are:



Limonene is an important volatile emitted by the holm oak (*Quercus ilex*), and acts as allelochemical in inhibiting seed germination of other plant species.

3.0.3 Bicyclic Monoterpenes



Pinene is, as limonene, an allelochemichal emitted by the roots of *Quercus ilex*. Camphor and pinene are also allelochemicals emitted by *Salvia leucophylla*.

any monoterpenes possess antitumor activity in animal and cell models. They have also antioxidant properties, -terpene and hydroxytyrosol being among the most effective.

3.1 Sesquiterpenes

Sesquiterpenoids are defined as the group of 15 carbon compounds derived by the assembly of 3 isoprenoid units and they are found mainly in higher plants but also in invertebrates. Sesquiterpenes, with monoterpenes, are an important constituent of essential oils in plants. They are the most diverse group of isoprenoids. In plants, they function as pheromones and juvenile hormones. Sesquiterpene structures present several acyclic, mono-, bi-, tri-, and tetracyclic systems.

3.1.1 Acyclic sesquiterpenes

The acyclic representative are also called farnesans, term derived from the basic structure, farnesol. Farnesol and nerolidol are very common and are isolated from essential oils of various sources.

Farnesol is widely distributed in many essential oils such as citronella, neroli, cyclamen, lemon grass, tuberose, rose, musk, and balsam. It is used in perfumery to emphasize the odors of perfumes. Moreover, it is a natural pesticide for mites and is also a pheromone for several insects and mammals, including elephants (teritorial marking, individual recognition, mate attraction).

Farnesene, an analogue of farnesol, is known to act as an alarm pheromone in aphids. Released during predator attack, it causes aphids to stop feeding, disperse, and give birth to winged (rather than wingless) forms, which leave their host plants.

Farnesene

Nerolidol is present in neroli, ginger, jasmine, lavender, tea tree and lemon grass. The aroma of nerolidol is woody and reminiscent of fresh bark. It is used as a flavoring agent and in perfumery. It was also shown to be produced by the leaves of a large number of plant species in response to herbivory insects and then to be transformed into a C11-homoterpene (4,8-dimethyl-1,3,7-nonatriene) which attracts predatory insects.

4,8-Dimethyl-1,3,7-nonatriene

3.1.2 Cyclic sesquiterpenes

Abscisic acid plays a key role in plants in the regulation of stomatal closure by regulating ion channel activities and water exchanges across the plasma membrane of guard cells.

Cyclic ADP-ribose (cADPR) has been shown to mediate signaling of abscisic acid in the drought-stress response leading to activation of gene transcription and to stomatal closure. Abscisic acid has also a variety of roles in plant development, bud and seed dormancy, germination, cell division and movement. It induces storage protein synthesis in seeds and may be involved in defense against insect attack.

Cadalene has the cadinane skeleton and is present in essential oils and in many plants. It is used as a biomarker in paleobotanic studies. In connection with retene (1-methyl-7-isopropyl phenanthrene), it enables the estimation in sediments of the importance of Pinaceae in ancient forests.

Some important Sesquiterpenes

Cadalene

Gossypol is a sesquiterpene dimer found in cotton that is formed from two cadinane units. All cotton plant contains gossypol. That terpene occurs as a mixture of two enantiomers but each has different biological activities. For non-ruminant animals (rodents, chickens, humans), (–)-gossypol is significantly more toxic than the (+) enantiomer. It has anticancer properties and inhibits male fertility in humans. In contrast, cotton plants containing high levels of (+)-gossypol are resistant to insect damage. These terpenes must be removed from the plant parts and oil before use as animal foods.

Capsidiol is a sesquiterpenoid compound that accumulates in tobacco *Nicotiana tabacum* and chili pepper *Capsicum annuum* in response to fungal infection. It is considered as a phytoalexin.

Capsidiol

3.2 Diterpenes

They have 20 carbon atoms and are derived from geranylgeraniol pyrophosphate. They are of fungal or plant origin and are found in resins, gummy exudates, and in the resinous high-boiling fractions remaining after distillation of essential oils. Diterpenoid groups that are physiologically active include: vitamin A activity (retinol), phytohormones that regulate plant growth and germination, e.g. gibberellins, fungal hormones that stimulate the switch from asexual to sexual reproduction, e.g. trisporic acid; disease resistance agents (phytoalexins), e.g. casbene and podocarpic acid, the anticancer drug,

taxol, from the bark of the yew tree, the cancer promoter, phorbol, and natural cannabinoids.

The diterpenes have exceptionally open chain, as found in geranylgeraniol or phytol which forms a part of chlorophyll and the side chain of vitamin E and K, and crocetin which is a diacid diterpenoid and the lipid part of the crocins, glycosylated derivatives present in saffron.

Examples of diterpene substances are given below:

Steviol is the aglycone of *stevia*'s sweet glycosides, one of them being formed by replacing one hydrogen atom (bottom) with glucose via an ester link, and another hydrogen atom (top) with a disaccharide (glucose and rhamnose). The steviol glycosides are responsible for the sweet taste of the leaves of the *stevia* plant (*Stevia rebaudiana*, Asteraceae). These compounds are 40 to 300 times sweeter than sucrose. They are developed to be used in sweet drinks.

Steviol

Retene is present in tars obtained by distillation of resinous wood, it is an important pollutant eliminated by the paper factories. This diterpene is present in geological sediment where it is formed by diagenesis from abietic acid, several intermediates having been recognized.

Retene

Dehydroleucodine was isolated from *Artemisia douglasiana*, a popular medicine in Argentina and was shown to have several physiological and therapeutic properties: anti-proliferative activity in G2 phase, cytoprotective agent for gastric ulcers and a general antioxidant.

Cafestol and kahweol are present in high concentrations (up to 18% diterpene esters) in the oil derived from coffee beans. The only difference between cafestol and kahweol is an extra double bond present in the second cycle of kahweol. These diterpenes are esterified with one fatty acid (C14 to C24), palmitic and linoleic acids being the major esterified fatty acids.

Phorbol is a diterpene isolated in 1934 from croton oil (seeds of *Croton tiglium*). Various fatty acid esters of phorbol have important biological properties, the most notable of which is the capacity to act as tumor promoters through activation of protein kinase C as they mimic diacylglycerols. The most common phorbol ester is phorbol-12-myristate-13-acetate (PMA), which is used as a research tool in models of carcinogenesis.

Phorbol

R: myristic acid (C14:0), Ac: acetate group

Cannabinoids are a group of diterpenes present in Cannabis (*Cannabis sativa L*). All these substances are structurally related to tetrahydrocannabinol (THC) and are able to bind to specific cannabinoid receptors.

Tetrahydrocannabinol (THC)

3.3 Sesterpenes

They are derived from geranylfarnesol pyrophosphate and have 25 carbon atoms. They were isolated from insect protective waxes and from fungal sources.

Three examples of sesterpenes are shown below.

Variously unsaturated and branched sesterpenes, known as Haslenes, were found in species of diatomaceous algae. They are widely distributed and abundant in marine sediments. Several haslenes were found to be produced by a iatom iatom *Haslea ostrearia* according to the culture temperature and were shown to have cytostatic. One of them is shown below.

3.4 Triterpenoids

They form a large group of natural substances which includes steroids and consequently sterols.

Squalene is the immediate biological precursor of all triterpenoids.

Squalene epoxide (2,3-oxidosqualene) is produced by the enzyme squalene epoxidase which use NADPH and oxygen to oxidize squalene. This metabolic step is the first in sterol biosynthesis leading to the formation of lanosterol or cycloartenol.

Squalane is a completely saturated derivative of squalene. Present in sebum, it is largely used as a component in many cosmetic products. It is obtained by hydrogenation of squalene extracted from olive oil. The

large groups of *steroids*, including sterols, are present in very small amounts in bacteria but at larger amounts in plants and animals while the hopanoids are very abundant in

prokaryotes where they replace cholesterol.

Among the large number of triterpenoid structures, some of them are shown below.

SELF-ASSESSMENT EXERCISE

- i. Highlight the five major classes of terpenes and give examples.
- ii. Describe sesquiterpenes and their biological importance.

4.0 CONCLUSION

Terpenes are hydrocarbons resulting from the combination of several isoprene units. Terpenoids can be thought of as modified terpenes, wherein methyl groups have been moved or removed, or oxygen atoms added. Just like terpenes, the terpenoids can be classified according to the number of isoprene units used. The classes of terpenes include monoterpenes, diterpenes, triterpenes, sesterpenes and sesquiterpenes.

5.0 SUMMARY

In summary, we have learnt that:

- i. Terpenes and terpenoids are so numerous and widespread in nature.
- ii. Terpenes are classified based on the number of their isoprene units.
- iii. The classes of terpenes include monoterpenes, diterpenes, triterpenes, sesterpenes and sesquiterpenes.

6.0 TUTOR-MARKED ASSIGNMENT

- 1. Distinguish between acyclic and cyclic sesquiterpenes
- 2. Name one example each of acyclic and cyclic sesquiterpenes and describe them
- 3. Name one example of natural source of acyclic and cyclic sesquiterpenes

7.0 REFERENCES/FURTHER READING

- Plemenkov VV (2001). Introduction to the Chemistry of Natural Compounds. Kazan.
- Paul M Dewick (2002). Medicinal Natural Products. A Biosynthetic Approach. Second Edition.

Wiley. ISBN 0471496405.

- Timothy J. Brocksom, Kleber T. de Oliveira and André L. Desiderá (2017): The Chemistry of the Sesquiterpene Alkaloids, J. Braz. 28 (6) http://dx.doi.org/10.21577/0103-5053.20170049
- Bahl, B. S. and Bahl, A..(2007). Advanced Organic Chemistry of Natural Product. Charp and Company Ltd.

MODULE 2 STEROIDS

UNIT 1 NATURE OF STEROIDS

CONTENTS

- 1.0 Introduction
- 2.0 Objectives
- 3.0 Definition of Steroid
 - 3.1 Structure of Steroid
 - 3.2 Classification
 - 3.2.1 Taxonomical/Functional Classification
 - 3.2.2 Structural Classification
 - 3.3 Metabolism of Steroid
 - 3.3.1 Steroid Synthesis
 - 3.3.2 Steroidogenesis
 - 3.3.3 Degradation and Elimination of Steroids
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 References/Further Reading

1.0 INTRODUCTION

The core of steroids is composed of twenty carbon atoms bonded together that take the form of four fused rings: three cyclohexane rings (designated as rings A, B, and C in the figure to the right) and one cyclopentane ring (the D ring). The steroids vary by the functional groups attached to this four ring core and by the oxidation state of the rings. Sterols are special forms of steroids, with a hydroxyl group at position-3 and a skeleton derived from cholestane.

2.0 OBJECTIVES

By the end of this unit, students should be able to:

- describe the structure of a steroid
- give various categories of steroids
- discuss the biosynthesis of steroids


3.0 MAIN CONTENT

Definition of Steroid

A steroid is a type of organic compound that contains a specific arrangement of four cycloalkane rings that are joined to each other. Examples of steroids include the dietary fat cholesterol, the sex hormones estradiol and testosterone, and the anti-inflammatory drug dexamethasone.

Hundreds of distinct steroids are found in plants, animals, and fungi. All steroids are made in cells either from the sterois lanosterol (animals and fungi) or from cycloartenol (plants). Both lanosterol and cycloartenol are derived from the cyclization of the triterpene squalene.

IUPAC recommended ring lettering (left) and atom numbering (right) of cholestane, a prototypical steroid skeleton. The four rings A-D form the gonane nucleus of the steroid.



26

Stick model of the steroid lanosterol. The total number of carbons (30) reflects its triterpenoid origin.

3.1 Structure of Steroid

Steroids are a class of organic compounds with a chemical structure that contains the core of gonane or a skeleton derived therefrom. Usually, methyl groups are present at the carbons C-10 and C-13. At carbon C-17 an alkyl side chain may also be present.

Numbering of carbon

atoms in gonane

The basic skeleton of a steroid, Cholestane, a typical steroid with standard stereo orientations.

R is a side-chain.

Gonane is the simplest possible steroid and is composed of seventeen carbon atoms, bonded together to form four fused rings. The three cyclohexane rings (designated as rings A, B, and C in the figure above right) form the skeleton of phenanthrene; ring D has a cyclopentane structure. Hence, together they are called cyclopentaphenanthrene.

Commonly, steroids have a methyl group at the carbons C-10 and C-13 and an alkyl side chain at carbon C-17. Further, they vary by the configuration of the side chain, the number of additional methyl groups and the functional groups attached to the rings. For example the hydroxyl group at position C-3 in sterols.

Cholesterol Cholic acid Medrogestone

3.2 Classification

3.2.1 Taxonomical/Functional Classification


Some of the common categories of steroids:

- 1. Animal steroids
 - a. Insect steroids: These include Ecdysteroids such as ecdysterone
 - b. Vertebrate steroids (steroid hormones)

Sex steroids are a subset of sex hormones that produce sex differences or support reproduction.

They include androgens, estrogens, and progestagens.

Corticosteroids include glucocorticoids and mineralocorticoids. Glucocorticoids regulate many aspects of metabolism and immune function, whereas mineralocorticoids help maintain blood volume and control renal excretion of electrolytes. Most medical 'steroid' drugs are corticosteroids.



Anabolic steroids are a class of steroids that interact with androgen receptors to increase muscle and bone synthesis. There are natural and synthetic anabolic steroids. In popular language, the word "steroids" usually refers to anabolic steroids.

Cholesterol, which modulates the fluidity of cell membranes and is the principal constituent of the plaques implicated in atherosclerosis.

Anabolic steroids

2. Plant steroids: Phytosterols, Brassinosteroids

3. Fungus steroids: Ergosterols

3.2.2 Structural Classification


This is based upon the chemical composition of the steroids. Examples from this classification include:

Class	Evennles	Number	of	carbon
	Examples	atoms		
Cholestane				
S	cholesterol	127		
	cholic			
Cholanes	acid	24		
	progesteron	1		
Pregnanes	e	21		
Androstanes	S			
testosterone		19		
Estranes	estradiol	18		

Gonane (or steroid nucleus) is the parent (17-carbon tetracyclic) hydrocarbon molecule without any alkyl sidechains.

3.3 Metabolism of Steroid

Steroids include estrogen, cortisol, progesterone, and testosterone. Estrogen and progesterone are made primarily in the ovary and in the placenta during pregnancy, and testosterone in the testes. Testosterone is also converted into estrogen to regulate the supply of each, in the bodies of both females and males. Certain neurons and glia in the central nervous system (CNS) express the enzymes that are required for the local synthesis of pregnane neurosteroids, either *de novo* or from peripherally-derived sources. The rate-limiting step of steroid synthesis is the conversion of cholesterol to pregnenolone, which occurs inside the mitochondrion.




Steroid metabolism is the complete set of chemical reactions in organisms that produce, modify, and consume steroids. These metabolic pathways include:

steroid synthesis – the manufacture of steroids from simpler precursors steroidogenesis – the interconversion of different types of steroids

steroid degradation.


3.3.1 Steroid Synthesis

Steroid biosynthesis is metabolic pathway that produces steroids from simple precursors. This pathway is carried out in different ways in animals than in many other organisms, making the pathway a common target for antibiotics and other anti-infective drugs. In addition, steroid metabolism in humans is the target of cholesterol-lowering drugs such as statins. It starts in the mevalonate pathway in humans, with Acetyl-CoA as building blocks, which form DMAPP and IPP. In following steps, DMAPP and IPP form lanosterol, the first steroid.



Simplified version of latter part of steroid synthesis pathway, where the intermediates isopentenyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP) form geranyl pyrophosphate (GPP), squalene and, finally, lanosterol, the first steroid in the pathways.

Mevalonate pathway



Mevalonate pathway

The mevalonate pathway or HMG-CoA reductase pathway starts with and ends with dimethylallyl pyrophosphate (DMAPP) and isopentenyl pyrophosphate (IPP).

Several key enzymes can be activated through DNA transcriptional regulation on activation of SREBP (Sterol Regulatory Element-Binding Protein-1 and -2). This intracellular sensor detects low cholesterol levels and stimulates endogenous production by the HMG-CoA reductase pathway, as well as increasing lipoprotein uptake by up-regulating the LDL receptor. Regulation of this pathway is also achieved by controlling the rate of translation of the mRNA, degradation of reductase and phosphorylation.

Pharmacology

A number of drugs target the *mevalonate pathway*:

Statins (used for elevated cholesterol levels)

Bisphosphonates (used in treatment of various bone-degenerative diseases)

Plants and bacteria

In plants and bacteria, the non-mevalonate pathway uses pyruvate and glyceraldehyde 3-phosphate as substrates.

DMAPP to lanosterol


Isopentenyl pyrophosphate and dimethylallyl pyrophosphate donate isoprene units, which are assembled and modified to form terpenes and isoprenoids, which are a large class of lipids that include the carotenoids, and form the largest class of plant natural products.

Here, the isoprene units are joined together to make squalene and then folded up and formed into a set of rings to make lanosterol. Lanosterol can then be converted into other steroids such as cholesterol and ergosterol.

3.3.2 Steroidogenesis

Steroidogenesis is the biological process by which steroids are generated from cholesterol and transformed into other steroids. The pathways of steroidogenesis differ between different species, but the pathways of human steroidogenesis are shown in the figure.

Lanosterol can then be converted into other steroids such as cholesterol and ergosterol.



Human Steroidogenesis

Products of steroidogenesis include:

- i. androgens: testosterone
- ii. estrogens and progesterone
- iii. corticoids: cortisol and aldosterone

3.3.3 Degradation and Elimination of Steroids

Steroids are oxidized mainly by cytochrome P450 oxidase enzymes, such as CYP3A4. These reactions introduce oxygen into the steroid ring and allow the structure to be broken up by other enzymes, to form bile acids as final products. These bile acids can then be eliminated through secretion from the liver in the bile. The expression of this oxidase gene can be up-regulated by the steroid sensor PXR when there is a high blood concentration of steroids.

SELF-ASSESSMENT EXERCISE

- i. What are steroids?
- ii. State the classes of steroids based on functions
- iii. Briefly describe steroidogenesis

4.0 CONCLUSION

Sterols are special forms of steroids, with a hydroxyl group at position-3 and a skeleton derived from cholestane. They can be classified based on their functions as well as their structures. Their metabolic pathways include steroid synthesis, steroidogenesis and steroid degradation.

5.0 SUMMARY

In summary, we have learnt that

- i. Steroids are a class of organic compounds with a chemical structure that contains the core of gonane or a skeleton derived therefrom.
- ii. Gonane is the simplest possible steroid and is composed of seventeen carbon atoms, bonded together to form four fused rings.
- iii. Steroids are produced in plants and animals as well as microorganism
- iv. Examples of steroids include oestrogen, progesterone, corticosterone, ergosterol and cholesterol

6.0 TUTOR-MARKED ASSIGNMENT

Describe the structure of gonane.

What is the biological importance of corticosteroids.

7.0 REFERENCES/FURTHER READING

- Paul M Dewick (2002). Medicinal Natural Products. A Biosynthetic Approach. Second Edition. Wiley. ISBN 0471496405.
- Plemenkov VV (2001). Introduction to the Chemistry of Natural Compounds. Kazan.
- Cfe Higher Chemistry Success Guide by Bob Wilson | Jun 15, 2015
- Finar, I. L.(2006). Stereochemistry and the chemistry of natural Product. Vol.2 (5thedn.), Pearson Education Ltd, Patparganj Delhi, India.
- Stephen P. S. (2006). Natural Products Chemistry at a Glance. Blackwell Publishers, UK.

MODULE 3 ALKALOIDS

UNIT 1 NATURE OF ALKALOIDS


CONTENTS

- 1.0 Introduction
- 2.0 Objectives
- 3.0 Definition of Alkaloid
 - 3.1 History of Alkaloid
 - 3.2 Properties of Alkaloid
 - 3.3 Distribution in Nature
 - 3.4 Extraction
 - 3.5 Biosynthesis of Alkaloids
 - 3.5.1 Synthesis of Schiff bases
 - 3.5.2 Mannich reaction
 - 3.5.3 Dimer alkaloids
 - 3.6 The biological role of Alkaloids
 - 3.7 Applications of Alkaloids
 - 3.7.1 In medicine
 - 3.7.2 In agriculture
 - 3.7.3 Use as psychoactive drugs
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 References/Further Reading

1.0 INTRODUCTION


Many alkaloids are toxic to other organisms. They often have pharmacological effects and are used as medications, as recreational drugs, or in entheogenic rituals. Examples are the local anesthetic and stimulant, cocaine; the stimulant, caffeine; nicotine; the analgesic, morphine; the antibacterial, berberine; the anticancer compound, vincrinstine; the antihypertension agent, resepine; the cholinomimeric, galatamine; the spasmolysis agent, atropine; the vasodilator, vincamine; the anti-arhythmia compound, quinidine; the anti-asthma therapeutic, ephedrine; and the antimalarial drug, quinine. Although alkaloids act on a diversity of metabolic systems in humans and other animals, they almost uniformly invoke a bitter taste.

nicotine



Vincristine

Galanthamine



Vincamine

2.0 OBJECTIVES

By the end of this unit, students should be able to:

- know what alkaloids are
- understand the properties of alkaloids
- Know modes of extraction of alkaloids
- comprehend the applications of alkaloids in human endeavour

3.0 MAIN CONTENT

Definition of Alkaloid

Alkaloid is a group of naturally occurring chemical compounds which mostly contain basic nitrogen atoms. This group also includes some related compounds with neutral and even weakly acidic properties. Also some synthetic compounds of similar structure are attributed to alkaloids. In addition to carbon, hydrogen and nitrogen, alkaloids may also contain oxygen, sulfur and more rarely other elements such as chlorine, bromine and phosphorus.


3.1 History of Alkaloid


Alkaloid-containing plants were used by humans since ancient times for therapeutic and recreational purposes. For example, medicinal plants have been known in the Mesopotamia at least around 2000 BC. The *Odyssey* of Homer referred to a gift given to Helen by the Egyptian queen, a drug bringing oblivion. It is believed that the gift was an opium-containing drug. A Chinese book on houseplants written in 1st-3rd centuries BC mentioned a medical use of Ephedra and opium poppies. Also, coca leaves were used by South American Indians since ancient times.

Studies of alkaloids began in the 19th century. In 1804, the German chemist Friedrich Sertürner isolated from opium a "soporific principle"


(Latin: *Principium somniferum*), which he called "Morphium" in honor of Morpheus, the Greek god of dreams; in German and some other Central-European languages, this is still the name of the drug. The term "morphine", used in English and French, was given by the French physicist Joseph Louis Gay-Lussac).

A significant contribution to the chemistry of alkaloids in the early years of its development was made by the French researchers Pierre Joseph Pelletier and Joseph Bienaimé Caventou who discovered quinine (1820) and strychnine (1818).






Several other alkaloids were discovered around that time, including xanthine (1817), atropine (1819), caffeine (1820), coniine (1827), nicotine (1828), colchicine (1833), sparteine (1851) and cocaine (1860).



Coniine

nicotine



Colchicines

The first complete synthesis of an alkaloid was achieved in 1886 by the German chemist Albert Ladenburg. He produced conine by reacting 2-methylpyridine with acetaldehyde and reducing the resulting 2-propenyl pyridine with sodium. The development of the chemistry of alkaloids

was accelerated by the emergence of spectroscopic and chromatographic methods in the 20th century, so that by 2008 more than 12,000 alkaloids had been identified.

3.2 Properties of Alkaloid

Most alkaloids contain oxygen; those compounds are usually colorless crystals at ambient conditions. Oxygen-free alkaloids, such as nicotine or coniine, are typically volatile, colorless, oily liquids. Some alkaloids are colored, like berberine (yellow) and sanguinarine (orange).

Most alkaloids are weak bases, but some are amphoteric, for example theobromine and theophylline. Most alkaloids are poorly soluble in water but readily dissolve in organic solvents, such as diethyl ether, chloroform and 1,2-dichloroethane. However, caffeine dissolves well in boiling water. With acids, alkaloids form salts of various strengths. Those salts are usually soluble in water and alcohol and poorly soluble in most organic solvents. Exceptions include scopolamine hydrobromide which is soluble in organic solvents and water-soluble quinine sulfate.

Most alkaloids have a bitter flavor. It is believed that plants evolved the ability to produce these bitter substances, many of which are poisonous, in order to protect themselves from animals; however, animals in turn evolved the ability to detoxify alkaloids. Some alkaloids can produce developmental defects in the offspring of animals that consume them but cannot detoxify them. A characteristic example is the alkaloid cyclopamine, which is present in the leaves of corn lily. During the 1950s, up to 25% lambs born by sheep that had grazed on corn lily suffered serious facial defects. Those defects ranged from deformed jaws to cyclopia. After decades of research, in 1980s, the substance that was responsible for the deformities was identified as the alkaloid 11-deoxyjervine, which was renamed cyclopamine.

3.3 Distribution in Nature

The alkaloid content in plants is usually within a few percent and is inhomogeneous over the plant tissues. Depending on the type of plants, the maximum concentration is observed in the leaves (black henbane), fruits or seeds (Strychnine tree), root (*Rauwolfia serpentina*) or bark (cinchona). Furthermore, different tissues of the same plants may contain different alkaloids.

Beside plants, alkaloids are found in certain types of fungi, such as psilocybin in the fungus of the genus Psilocybe, and in animals, such as bufotenin in the skin of some toads. Many marine organisms also contain alkaloids. Some amines, such as adrenaline and serotonin, which


play an important role in higher animals, are similar to alkaloids in their structure and biosynthesis and are sometimes called alkaloids.

3.4 Extraction


Because of the structural diversity of alkaloids, there is no single method of their extraction from natural raw materials. Most methods exploit the property of most alkaloids to be soluble in organic solvents but not in water, and the opposite tendency of their salts Scheme (1 and 2).

Most plants contain several alkaloids. Their mixture is extracted first and then individual alkaloids are separated. Plants are thoroughly ground before extraction. Most alkaloids are present in the raw plants in the form of salts of organic acids. The extracted alkaloids may remain as salts or change into bases. Base extraction is achieved by processing the raw material with alkaline solutions and extracting the alkaloid bases with organic solvents, such as 1,2-dichloroethane, chloroform, diethyl ether or benzene. Then, the impurities are dissolved by weak acids; this converts alkaloid bases into salts which are washed away with water. If necessary, an aqueous solution of alkaloid salts is again made alkaline and treated with an organic solvent. The process is repeated until the desired purity is achieved.

Scheme 1 illustrate soxhlet extraction using methanol and Scheme 2 illustrate soxhlet extraction using n-hexane



Scheme 1: Extraction and Isolation of Alkaloids from Plant Material Using Methanol as Solvent



Scheme 2: Extraction and Isolation of Alkaloids from Plant Material Using n- Hexane as Solvent

In the acidic extraction, the raw plant material is processed by a weak acidic solution (e.g., acetic acid in water, ethanol or methanol). A base is then added to convert alkaloids to basic forms which are extracted with organic solvent (if the extraction was performed with alcohol, it is removed first, and the remainder is dissolved in water). The solution is purified as described above.

Alkaloids are separated from their mixture using their different solubility in certain solvents and different reactivity with certain reagents or by distillation.

3.5 Biosynthesis of Alkaloids

Biological precursors of most alkaloids are amino acids, such as ornithine, lysine, phenylalanine, tyrosine, tryptophan, histidine, aspartic acid and anthranilic acid. Nicotinic acid can be synthesized from tryptophan or aspartic acid. Ways of alkaloid biosynthesis are too numerous and cannot be easily classified. However, there are a few typical reactions involved in the biosynthesis of various classes of alkaloids, including synthesis of Schiff bases and Mannich reaction.

3.5.1 Synthesis of Schiff bases

Schiff bases can be obtained by reacting amines with ketones or aldehydes. These reactions are a common method of producing C=N bonds.

$$\begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c} R \\ R_1 \end{array} \longrightarrow \begin{array}{c} R \\ R_2 \end{array} \longrightarrow \begin{array}{c}$$

In the biosynthesis of alkaloids, such reactions may take place within a molecule, such as in the synthesis of piperidine:

3.5.2 Mannich reaction

An integral component of the Mannich reaction, in addition to an amine and a carbonyl compound, is a carbanion, which plays the role of the nucleophile in the nucleophilic addition to the ion formed by the reaction of the amine and the carbonyl.

The Mannich reaction can proceed both intermolecularly and intramolecularly:

$$\longrightarrow$$

3.5.3 Dimer alkaloids

In addition to the described above monomeric alkaloids, there are also dimeric, and even trimeric and tetrameric alkaloids formed upon condensation of two, three and four monomeric alkaloids. Dimeric alkaloids are usually formed from monomers of the same type through the following mechanisms:

Mannich reaction, resulting in, e.g., voacamine Michael reaction (villalstonine).

Condensation of aldehydes with amines (toxiferine).

Oxidative addition of phenols (dauricine, tubocurarine). Lactonization (carpaine).

Voacamine

Toxiferine

Dauricine

Tubocurarine

Carpaine

3.6 The biological role of Alkaloids

The role of alkaloids for living organisms which produce them is still unclear. Initially, it was assumed that the alkaloids are the final products of nitrogen metabolism in plants, as urea in mammals. Most of the known functions of alkaloids are related to protection. For example, aporphine alkaloid liriodenine produced by the tulip tree protects it from parasitic mushrooms. In addition, presence of alkaloids in the plant prevents insects and chordate animals from eating it. However, some animals adapted to alkaloids and even use them in their own metabolism. Such alkaloid-related substances as serotonin, dopamine and histamine are important neurotransmitters in animals. Alkaloids are also known to regulate plant growth.

3.7 Applications of Alkaloids

3.7.1 In medicine

Medical use of alkaloid plants has a long history, and thus when the first alkaloids were synthesized in the 19th century, they immediately found application in clinical practice. Many alkaloids are still used in medicine, usually in the form of salts, including the following:

Table 3.0: Medical uses of alkaloids

Alkaloid	Action		
Ajmaline	Antiarrhythmic		
Atronina			
Atropine,	Antichalinancia		
scopolamine	Anticholinergic		
Vinblastine,			
vincristine	Antitumor		
Vincamin			
e	vasodilating, antihypertensive		
	cough		
Codeine	medicine		
<u> </u>	anestheti		
Cocaine			
Colchicin	<u>c</u>		
	mamady for cout		
<u>e</u>	remedy for gout		
Morphine	analgesic		
Reserpin	antihypertensiv		
e	e		
Tubocurarin			
e	muscle relaxant		
Physostigmin	inhibitor of		
e	acetylcholinesterase		
Quinidin			
e	Antiarrhythmic		
Quinin	,		
e	antipyretics, antimalarial		
Emetin			
	antiprotozoal		
<u>e</u>	agent		



Many synthetic and semisynthetic drugs are structural modifications of the alkaloids, which were designed to enhance or change the primary effect of the drug and reduce unwanted side effects.

For example, naloxone, an opioid receptor antagonist, is a derivative of the baine which is present in opium.

Thebaine

Naloxone

3.7.2 In agriculture

Prior to the development of a wide range of relatively low-toxic synthetic pesticides, some alkaloids, such as salts of nicotine and anabasine, were used as insecticides. Their use was limited by their high toxicity to humans.

3.7.3 Use as psychoactive drugs

Preparations of plants containing alkaloids and their extracts, and later pure alkaloids have long been used as psychoactive substances. Cocaine and cathinone are stimulants of the central nervous system. Mescaline and many of indole alkaloids (such as psilocybin, dimethyltryptamine and ibogaine) have hallucinogenic effect. Morphine and codeine are strong narcotic pain killers.

There are alkaloids that do not have strong psychoactive effect themselves, but are precursors for semi-synthetic psychoactive drugs.

For example, ephedrine and pseudoephedrine are used to produce methcathinone and methamphetamine.

SELF-ASSESSMENT EXERCISE

- i. In tabular form, state 10 examples of alkaloids and their therapeutic importance.
- ii. Describe the modes of extraction of alkaloids.
- iii. With examples, describe the properties of alkaloids.

4.0 CONCLUSION

Alkaloids are produced by a large variety of organisms, including bacteria, fungi, plants, and animals and are part of the group of natural products (also called secondary metabolites). Many alkaloids can be purified from crude extracts by acid-base extraction. The boundary between alkaloids and other nitrogen-containing natural compounds is not clear-cut. Compounds like amino acid peptides, proteins, nucleotides, nucleic acid, amines and antibiotics are usually not called alkaloids. Natural compounds containing nitrogen in the exocyclic position (mescaline, serotonin, dopamine, etc.) are usually attributed to amines rather than alkaloids.

5.0 SUMMARY

- i. Alkaloid is a group of naturally occurring chemical compounds which mostly contain basic nitrogen atoms.
- ii. Most alkaloids are weak bases, but some are amphoteric, for example theobromine and theophylline.
- iii. Alkaloids are found mostly in plants, marine organisms and in some fungi such as genus Psilosybe.
- iv. Alkaloids are of widespread importance especially in medicine and agriculture.

6.0 TUTOR-MARKED ASSIGNMENT

- 1. List five properties of alkaloids
- 2. How is alkaloid distributed in nature?

7.0 REFERENCES/FURTHER READING

Tadeusz, A. (2007): Alkaloids - secrets of life. Amsterdam: Elsevier.

Fattorusso, E and Taglialatela-Scafati, O. (2008): *Modern Alkaloids: Structure, Isolation, Synthesis and Biology*. Wiley-VCH.

- Manfred, H. (2002): *Alkaloids: Nature's Curse or Blessing?*. Wiley-VCH.
- Orekhov, A. P. (1955): *Chemistry alkaloids*, Academic Publisher (2nd ed.). USSR.
- Cfe Higher Chemistry Student Book by Tom Speirs | Nov 19, 2014
- Patrick GL (2013). "Medical Folklore". *An introduction to medicinal chemistry* (Fifth ed.). Oxford: Oxford University Press. ISBN 978-0-19-969739-7.

UNIT 2 CLASSIFICATION OF ALKALOIDS

CONTENTS

- 1.0 Introduction
- 2.0 Objectives
- 3.0 Classes of Alkaloids
 - 3.1 Major Classes of Alkaloids
 - 3.2 Main Classes of Alkaloids
- 4.0 Conclusion
- 5.0 Summary
- 6.0 Tutor-Marked Assignment
- 7.0 Refeferences/Further Reading

1.0 INTRODUCTION

Compared with most other classes of natural compounds, alkaloids are characterized by a great structural diversity and there is no uniform classification of alkaloids. Historically, first classification methods combined alkaloids by the common natural source, e.g., a certain type of plants. This classification was justified by the lack of knowledge about the chemical structure of alkaloids and is now considered obsolete.

2.0 OBJECTIVES

By the end of this unit, students should be able to:

- understand the diversity of alkaloids
- know the major classes of alkaloids
- examples of alkaloids in each group

3.0 MAIN CONTENT Classes of Alkaloids

3.1 Major Classes of Alkaloids

Alkaloids are often divided into the following major groups:

1. "True alkaloids", which contain nitrogen in the heterocycle and originate from amino acids. Their characteristic examples are atropine, nicotine and morphine. This group also includes some alkaloids which beside nitrogen heterocycle contain terpene (e.g. evonine) or peptide fragments (e.g. ergotamine). This group also includes piperidine alkaloids coniine and coniceine although they do not originate from amino acids.

- 2. "Protoalkaloids", which contain nitrogen and also originate from amino acids. Examples include mescaline, adrenaline and ephedrine.
- 3. Polyamine alkaloids – derivatives of putrescine, spermidine and spermine.
- 4. Peptide and cyclopeptide alkaloids.
- 5. Pseudalkaloids – alkaloid-like compounds which do not originate from amino acids. This group includes, terpene-like and steroidlike alkaloids, as well as purine-like alkaloids such as caffeine, theobromine and theophylline. Some authors classify pseudoalkaloids such compounds such as ephedrine and cathinone. Those originate from the amino acid phenylalanine, but acquire their nitrogen atom not from the amino acid but through transamination.

Some alkaloids do not have the carbon skeleton characteristic of their group. So, galantamine and homoaporphines do not contain isoquinoline fragment, but are generally attributed to isoquinoline alkaloids.

3.2 **Main Classes of Alkaloids**

Alkaloids with nitrogen heterocycles alkaloids)

Class

Pyrrolidine derivatives^[41]

Tropane derivatives^[44]

Major groups

Main synthesis steps Examples

Ornithine or arginine Cuscohygrine, putrescine methylputrescine hygrine, Nhygroline, Stachydrine

methyl- ¹-pyrroline

Atropine group Atropine,

Substitution in positions 3,

6 Ornithine or arginine scopolamine,

or 7 putrescine N- Hyoscyamine

methylputrescine

Cocaine group N-

Cocaine,

Substitution in positions 2 and methyl- 1-pyrroline ecgonine

Pyrrolizidine derivatives



Piperidine derivatives



Quinolizidine derivatives

Non-esters

Complex esters of monocarboxylic acids

Macrocyclic diesters

1-aminopyrrolizidines (lolines)

Lupinine group

Retronecine,

heliotridine,

In plants: ornithine

or laburnine

arginine

putrescine Indicine,

homospermidine lindelophin, retronecine [42] Sarracine

Platyphylline, Trichodesmine

In fungi: L-proline +

L-

homoserine N-(3- Loline, N-

amino-3- formylloline, *N*-carboxypropyl)prolineAcetylloline

norloline

Sedamine,

Lysine cadaverine lobeline,

anaferine,

¹-piperideine piperine

Octanoic acid Coniine,

coniceine

coniceine coniine

Lysine cadaverine Lupinine, ¹-piperideine Nupharidin

Ind	1.	•	٠.		
Ind	ΔI_{11}	711	11	n	0



derivatives



Cytisine group

Sparteine group Matrine group Ormosanine group

Lysine

semialdehyde of -

aminoadipic acid

pipecolic acid 1 indolizidinone Cytisine

Sparteine, lupanine, anahygrine

Matrine, oxymatrine, allomatridine

Ormosanine, piptantine

Swainsonine, castanospermine

Pyridine derivatives



Isoquinoline derivatives and related alkaloids [69]



Simple derivatives of pyridine

Polycyclic noncondensing pyridine derivatives

Polycyclic condensed pyridine derivatives

Sesquiterpene pyridine derivatives

Simple derivatives of isoquinoline

Derivatives of 1- and 3-isoquinolines

Derivatives of 1- and 4-phenyltetrahydroisoquinolines

Derivatives of 5-naftil-isoquinoline

Derivatives of 1- and 2-benzyl-izoquinolines

Cularine group

Pavines and isopavines

Benzopyrrocolines

Protoberberines

Trigonelline,

ricinine, arecoline

Nicotine,

Nicotinic acid nornicotine, dihydronicotinic acid anabasine, 1,2-dihydropyridine anatabine

Actinidine, gentianine, Pediculinine Evonine,

Nicotinic acid,

hippocrateine,

Isoleucine

Triptonine Salsoline, Lophocerine

N-

methylcoridaldine,

Noroxyhydrastinine

Cryptostilin

Tyrosine or

Ancistrocladine

phenylalanine

dopamine or

tyramine Papaverine,

(for alkaloids Amarillis) laudanosine, sendaverine

Cularine, yagonine

Argemonine,

amurensin

Cryptaustoline

Berberine,

Canadine,

Oxazole

derivatives^[92]

Phtalidisoquinolines

Spirobenzylisoquinolines

Ipecacuanha alkaloids Benzophenanthridines

Aporphines

Proaporphines

Homoaporphines

Homoproaporphines

Morphines

Homomorphines

Tropoloisoquinolines

Azofluoranthenes

Amaryllis alkaloids

Erythrite alkaloids
Phenanthrene derivatives
Protopins
Aristolactam
ophiocarpine,
mecambridine, corydaline
Hydrastine,
narcotine (Noscapine)
Fumaricine
Emetine,
protoemetine, ipecoside
Sanguinarine,
oxynitidine, corynoloxine
Glaucine, coridine,
liriodenine
Pronuciferine,
glaziovine
Kreysiginine,
multifloramine
Bulbocodine
Morphine, codeine,



thebaine, sinomenine

Kreysiginine,

androcymbine

Imerubrine

Rufescine,

imeluteine

Lycorine,

ambelline, tazettine, galantamine, montanine

Erysodine,

erythroidine

Atherosperminine

Protopine,

oxomuramine, corycavidine

Doriflavin

Annuloline,

Tyrosine tyramine halfordinol, texaline, texamine

Isoxazole derivatives



Thiazole derivatives



Quinazoline derivatives



Acridine derivatives



3,4-Dihydro-4-quinazolone derivatives

1,4-Dihydro-4-quinazolone derivatives

Pyrrolidine and piperidine quinazoline derivatives

Ibotenic acid Muscimol

1-Deoxy-D-xylulose 5-phosphate (DOXP), tyrosine, cysteine

Anthranilic acid or phenylalanine or ornithine

Anthranilic acid

Ibotenic acid, Muscimol

Nostocyclamide, thiostreptone

Febrifugine

Glycorine, arborine, glycosminine

Vazicine (peganine)

Rutacridone, acronicine

Quinoline derivatives



Simple derivatives of quinoline derivatives of 2 - quinolones and 4-quinolone

Tricyclic terpenoids

Furanoquinoline derivatives

Quinines

Anthranilic acid 3-carboxyquinoline

Tryptophan tryptamine strictosidine (with secologanin) korinanteal cinhoninonCusparine, echinopsine, evocarpine

Flindersine Dictamnine, fagarine, skimmianine Quinine quinidine cinchonine, cinhonidine

Indole derivatives

Non-isoprene indole alkaloids

Tryptophan
Simple indole derivatives
tryptamine or 5Serotonin,
psilocybin,

Imidazole derivatives



hydroxitriptofan

dimethyltryptamine (DMT), bufotenin

Harman, harmine, Simple derivatives of -

harmaline, carboline eleagnine

Physostigmine

(eserine),

Pyrroloindole alkaloids

physovenine, eptastigmine

etheramine,

Semiterpenoid indole alkaloids

Tryptophan chanoclavine

Ergotamine,

agroclavine

Ergot alkaloids

ergobasine,

elimoclavine

Ergosine

paspalic acid lysergic acid

Monoterpenoid indole alkaloids

Ajmalicine, sarpagine, vobasine, ajmaline, yohimbine, reserpine,

Corynanthe type alkaloids

mitragynine, group

Tryptophan strychnine and

(Strychnine

tryptamine strictosidine (with

secologanin)

brucine, aquamicine, vomicine)

Ibogamine, ibogaine, Voacangine

Vincamine,

Aspidosperma-type

Iboga-type alkaloids

alkaloids vincotine,

Aspidospermine

Histamine,
Directly from histidine pilocarpine,
pilosine, stevensine

Purine derivatives

N N

Xanthosine (formed in purine biosynthesis) 7 methylxantosine 7methyl xanthine theobromine

Caffeine theobromine theophylline saxitoxin caffeine

Alkaloids with nitrogen in the side chain (protoalkaloids)

-Phenylethylamine

derivatives

Tyrosine or

phenylalanine Tyramine,

dioxyphenilalanine ephedrine, dopamine adrenaline pseudoephedrine, and mescaline tyrosine mescaline,

tyramine cathinone,

phenylalanine 1- catecholamines phenylpropane-1,2- (adrenaline, dione cathinone noradrenaline,

ephedrine and dopamine) pseudoephedrine

Colchicine

alkaloids

Muscarine

Benzylamine

Putrescine

derivatives

Spermidine

derivatives

Spermine

derivatives

Tyrosine or phenylalanine

Colchicine,

dopamine

Colchamine

autumnaline Colchicines

Glutamic acid 3ketoglutamic acid muscarine (with pyruvic acid)

Muscarine, allomuscarine, epimuscarine, Epiallomuscarine

Phenylalanine with

Capsaicin,

valine, leucine or

dihydrocapsaicin, Nordihydrocapsaici

Isoleucine

Polyamines alkaloids

Paucine

ornithine putrescine spermidine

Lunarine,

Spermine

Codonocarpine

Verbascenine, Aphelandrine



Peptide alkaloids with a 13-membered cycle

Peptide alkaloids with a 14-membered cycle

Peptide alkaloids with a 15-membered cycle

Diterpenes

Steroids

4.0 CONCLUSION

Peptide (cyclopeptide) alkaloids

Numularine C, Numularine C type numularine S Ziziphin A, Ziziphin type sativanine H Frangulanine, Frangulanine type scutianine J Scutianine A Scutianine A type From different amino Integerrine, Integerrine type Acids discarine D Amphibine F, Amphibine F type spinanine A Amphibine B, Amfibine B type lotusine C

Mucronine A

Mucronine A type

Pseudoalkaloids (terpenes and steroids)

Mevalonic acid izopentenilpyrophosfate Aconitine, Licoctonine type geranyl pyrophosphate

delphinine

Solasodine,

Cholesterol, arginine solanidine, veralkamine

Recent classification of alkaloids are based on similarity of the carbon skeleton (e.g., indole, isoquinoline and pyridine-like) or biogenetic precursor (ornithine, lysine, tyrosine, tryptophan, etc.). However, they require compromises in borderline cases; for example, nicotine contains a pyridine fragment from nicotinamide and pyrrolidine part from ornithine and therefore can be assigned to both classes.

5.0 SUMMARY

There are five classes of alkaloids based on functions. These are:

- 1. "Protoalkaloids", which contain nitrogen and also originate from amino acids. Examples include mescaline, adrenaline and ephedrine.
- 2. Polyamine alkaloids derivatives of putrescine, spermidine and spermine.
- 3. Peptide and cyclopeptide alkaloids.
 - 4. Pseudalkaloids alkaloid-like compounds which do not originate from amino acids. E.g caffeine, theobromine and theophylline.

6.0 TUTOR-MARKED ASSIGNMENT

- 1. Highlight the five major classes of alkaloids
- 2. Describe with examples the true alkaloids

7.0 REFERENCES/FURTHER READING

Tadeusz, A. (2007): Alkaloids - secrets of life. Amsterdam: Elsevier.

Fattorusso, E and Taglialatela-Scafati, O. (2008): *Modern Alkaloids: Structure, Isolation, Synthesis and Biology*. Wiley-VCH.

Manfred, H. (2002): *Alkaloids: Nature's Curse or Blessing?*. Wiley-VCH.

- Orekhov, A. P. (1955): *Chemistry alkaloids*, Academic Publisher (2nd ed.). USSR.
- Cfe Higher Chemistry Student Book by Tom Speirs | Nov 19, 2014
- Patrick GL (2013). "Medical Folklore". *An introduction to medicinal chemistry* (Fifth ed.). Oxford: Oxford University Press. ISBN 978-0-19-969739-7.
- Tadeusz, A. (2007): Alkaloids secrets of life. Amsterdam: Elsevier.
- Fattorusso, E. and Taglialatela-Scafati, O. (2008): *Modern Alkaloids: Structure, Isolation, Synthesis and Biology*. Wiley-VCH.
- Manfred, H. (2002): Alkaloids: Nature's Curse or Blessing?. Wiley-VCH.
- Orekhov, A. P. (1955): Chemistry alkaloids (Acad. 2 ed.). M.: USSR.
- Stephen P. S. (2006). Natural Products Chemistry at a Glance. Blackwell Publishers ,UK.
- Higher Chemistry for Cfe. by John Anderson, Eric Allan, John Harris by John Anderson | Mar 1, 2012